

Finite Groups with Small Sums of Degrees of Some Non-linear Irreducible Characters

YAKOV BERKOVICH*

*Department of Mathematics and Computer Science, Research Institute of Afula,
University of Haifa, 31905 Haifa, Israel*

Communicated by Walter Feit

Received July 1993

IN MEMORY OF PROFESSOR S. A. AMITSUR

We obtain lower estimates for sums of degrees of some non-linear irreducible characters of non- p -nilpotent groups, and investigate the structure of groups for which these estimates are attained. © 1995 Academic Press, Inc.

Let p be a prime divisor of the order of a finite group G , P a fixed Sylow p -subgroup of G , $\text{Irr}(G)$ the set of all ordinary irreducible characters of G , and $\text{Irr}_1(G)$ the subset of all non-linear characters of $\text{Irr}(G)$. A group G is said to be p -nilpotent if it has a normal p -complement. Let

$$\begin{aligned}\text{Irr}_1(G, p') &= \{\chi \in \text{Irr}_1(G) \mid p \nmid \chi(1)\}, \\ \text{Irr}_1(G, P, p') &= \{\chi \in \text{Irr}_1(G, p') \mid p \mid |G/\ker \chi|\}, \\ \text{Irr}_1^0(G, p') &= \{\chi \in \text{Irr}_1(G, p') \mid G/\ker \chi \text{ is not } p\text{-nilpotent}\}, \\ \tau_1(G, p') &= \sum \chi, \quad \text{where } \chi \text{ runs over the set } \text{Irr}_1(G, p'), \\ T_1(G, p') &= \tau_1(G, p')(1).\end{aligned}$$

If we take, in the last two definitions, $\text{Irr}_1(G, P, p')$ or $\text{Irr}_1^0(G, p')$ instead of $\text{Irr}_1(G, p')$, we obtain the definitions of

$$\tau_1(G, P, p'), \quad T_1(G, P, p')(\tau_1^0(G, p'), \quad T_1^0(G, p')).$$

In this paper we obtain lower estimates of the numbers defined above and investigate the groups for which these estimates are attained.

Thompson (see [1s, Corollary 12.2]) showed that if $T_1(G, p') = 0$ then G is p -nilpotent. Our results generalize Thompson's theorem. Proposition 9 contains a new property of Thompson's groups. Moreover, as follows from this proposition (see Remark 1 following it), Thompson's groups are

*Supported in part by the Rashi Foundation and the Ministry of absorption of Israel.

solvable. The proof of this fact uses the classification of finite simple groups.

Let G' be the commutator subgroup of G , $\Phi(G)$ the Frattini subgroup of G , $Z(G)$ the centre of G , (A, B) a Frobenius group with kernel B and a complement A , $C(m)$ a cyclic group of order m , and $E(p^m)$ an elementary abelian group of order p^m . Denote by $\text{Irr}(\chi)$ the set of all irreducible constituents of a character χ . Let $\text{Lin}(G)$ be the set of all linear characters of G . In the sequel P denotes a fixed Sylow p -subgroup of G . If X is a set of characters of a group then $X^\# = X - \{\varepsilon\}$, where ε is the principal character of this group. Next, $O^p(G)$ denotes the subgroup generated by all p' -elements of G .

1

In this section we prove some introductory lemmas.

LEMMA 1. *Suppose that P' is normal in G . Then G is p -nilpotent if and only if G/P' is p -nilpotent.*

This is obvious since $P' \leq \Phi(G)$.

LEMMA 2. *Suppose that P is normal in G . Then*

$$T_1(G, P, p') = T_1(G/P', P/P', p').$$

Proof. Since $\text{Irr}_1(G/P', P/P', p') \subseteq \text{Irr}_1(G, P, p')$, it is sufficient to prove the inverse inclusion. Take $\chi = \text{Irr}_1(G, P, p')$, $\lambda \in \text{Irr}(\chi_P)$. Since $p \nmid \chi(1)$ and $\lambda(1) \mid \chi(1)$, $p \nmid \lambda(1)$ and $\lambda \in \text{Lin}(P)$, and hence $P' \leq \ker \lambda$ and $P' \leq \ker \chi$ (since λ is an arbitrary irreducible constituent of χ). Therefore χ is contained in $\text{Irr}(G/P', P/P', p')$. ■

LEMMA 3. *Suppose that P is normal in a non- p -nilpotent group G . If $p^\alpha \mid |G : G'|$ but $p^{\alpha+1} \nmid |G : G'|$, then*

- (a) $T_1(G, P, p') \geq |P : P'| - p^\alpha$.
- (b) $T_1(G, P, p') = |P : P'| - p^\alpha$ if and only if

$$G/P' = (H, P_1/P') \times P_2/P', |P_2/P'| = p^\alpha;$$

here $P_1 \in \text{Syl}_p(G')$.

Proof. In view of Lemmas 1 and 2 we may assume $P' = 1$, i.e., P is abelian.

(a, i) Assume that $\alpha = 0$, i.e., $P \leq G'$. Take $\lambda \in \text{Lin}^\#(P)$. By virtue of $P \leq G'$ one has $\text{Irr}(\lambda^{G'}) \subseteq \text{Irr}_1(G)$. Now $\text{Irr}_1(G) = \text{Irr}_1(G, p')$ by Ito's theorem on degrees [Is, Theorem 6.15]. Take $\chi \in \text{Irr}(\lambda^{G'})$. Then $P \cap \ker \chi = \ker \chi_P \leq \ker \lambda$ by reciprocity, so P is not contained in $\ker \chi$. Hence $\chi \in \text{Irr}_1(G, P, p')$. By reciprocity

$$\begin{aligned} \text{Lin}^\#(P) &\subseteq \text{Irr}(\tau_1(G, P, p)_P), \\ |P| - 1 &= |\text{Lin}^\#(P)| \leq T_1(G, P, p') \end{aligned}$$

and (a) is true for $\alpha = 0$.

(a, ii) Assume $\alpha > 0$. Take a p' -Hall subgroup H of G (Schur–Zassenhaus). Since HG' is normal in G one obtains $G = N_G(H)HG' = N_G(H)G'$ (the Frattini lemma and the Schur–Zassenhaus theorem). Now $N_G(H) = H \times P_2$, where $P_2 = P \cap N_G(H)$. In view of $C_G(P_2) \geq \langle P, H \rangle = G$ one has $P_2 \leq Z(G)$. Since P is abelian, $P \cap G' \cap Z(G) = 1$ (the transfer theorem), and we obtain

$$P_2 \cap G' = 1, \quad G = HG' \times P_2, \quad |P_2| = p^\alpha.$$

Take $P_1 \in \text{Syl}_p(G')$. In view of commutativity of P one has $O^{p'}(HG') = HG'$, so $P_1 \leq (HG')'$. Hence (a, i) implies $T_1(HG', P_1, p') \geq |P_1| - 1$. If $\lambda \in \text{Irr}_1(HG', P_1, p')$, $\mu \in \text{Lin}(P_2)$, then $\lambda \times \mu \in \text{Irr}_1(G, P, p')$, implying that

$$T_1(G, P, p') \geq T_1(HG', P_1, p')|P_2| \geq (|P_1| - 1)|P_2| = |P| - p^\alpha$$

and (a) is proved.

(b, i) Assume that $\alpha = 0$, i.e., $P \leq G'$. By this assumption $T_1(G, P, p') = |P| - 1$. It follows from the proof of (a, i) that $\lambda^{G'} \in \text{Irr}(G)$ for any $\lambda \in \text{Lin}^\#(P)$. So [K, Corollary 37.5.4] one has $G = (H, P)$. The inverse follows from the description of irreducible characters of Frobenius groups [Is, Theorem 6.34].

(b, ii) Assume $\alpha > 0$. As in (a, ii) one has $G = HG' \times P_2$, $|P_2| = p^\alpha$. Let $P_1 \in \text{Syl}_p(G')$. It follows from (a) and $(|P_1| - 1)|P_2| \leq T(HG', P, p')|P_2| \leq T(G, P, p') = |P| - p^\alpha$ that $T_1(HG', P_1, p') = |P_1| - 1$. Then (b, i) implies $HG' = (H, G')$. On the other hand, if G has the structure described above, then $T_1(G, P, p') = |P| - p^\alpha$. ■

LEMMA 4. Suppose that $P < H < G$. Then

- (a) $T_1(H, P, p') \leq T_1(G, P, p')$.
- (b) $T_1(H, p') \leq T_1(G, p')$.

Proof. (a) Take $\lambda \in \text{Irr}_1(H, P, p')$. Since $p \nmid \lambda^G(1) = |G : H|\lambda(1)$ there is $\chi \in \text{Irr}(\lambda^G)$ with $p \nmid \chi(1)$. By reciprocity $\chi(1) \geq \lambda(1)$, $H \cap \ker \chi \leq \ker \lambda$. Since P is not contained in $\ker \lambda$ it is also not contained in $\ker \chi$ and $\chi \in \text{Irr}_1(G, P, p')$. By reciprocity

$$\text{Irr}_1(H, P, p') \subseteq \text{Irr}(\tau_1(G, P, p')_H)$$

and the result follows.

(b) is proved similarly. ■

LEMMA 5. If G is not p -nilpotent then $T_1^0(G, p') \geq p - 1$.

Proof. If M is normal in G then $T_1^0(G, p') \geq T_1^0(G/M, p')$. Hence we may assume that all proper epimorphic images of G are p -nilpotent. Then G contains only one minimal normal subgroup M . Since G/M is p -nilpotent, $p \mid |M|$. Now $M \leq G'$ as the unique minimal normal subgroup of a non-abelian group G . Let us consider the following two cases.

(i) $p \nmid |G/M|$.

Then $P \leq M$. If $\chi \in \text{Irr}_1(G, P, p')$ then P and M are not contained in $\ker \chi$, so that χ is faithful and $G/\ker \chi = G$ is not p -nilpotent. Hence $\text{Irr}_1^0(G, p') = \text{Irr}_1(G, P, p')$. If $\lambda \in \text{Lin}^\#(P)$ then by virtue of $p \nmid \lambda^G(1)$ there is $\chi \in \text{Irr}(\lambda^G)$ with $p \nmid \chi(1)$. Obviously

$$\chi \in \text{Irr}_1(G, P, p') = \text{Irr}_1^0(G, p'),$$

so, by reciprocity, one obtains

$$T_1^0(G, p') \geq |\text{Lin}^\#(P)| = |P : P'| - 1 \geq p - 1.$$

(ii) $p \mid |G : M|$.

Let H/M be a normal p -complement of G/M . Since G/H is a p -group, H is not p -nilpotent. By Tate's theorem [Is, Theorem 6.31] the intersection $H \cap P = P_1$ is not contained in P' . Hence there is $\lambda \in \text{Lin}(P)$ such that P_1 is not contained in $\ker \lambda$. Take $\chi \in \text{Irr}(\lambda^G)$. Since $P_1 \leq G'$ and P_1 is not contained in $\ker \chi$ by reciprocity we have $\chi(1) > 1$. Since $p \nmid \lambda^G(1)$ we may choose χ so that $p \nmid \chi(1)$. Since P_1 is not contained in $\ker \chi$, M is not contained in $\ker \chi$, and χ is faithful. Since $G/\ker \chi = G$ is not p -nilpotent, $\chi \in \text{Irr}_1^0(G, p')$. As λ is an arbitrary character from $\text{Lin}(P/P') = \text{Lin}(P/P_1P')$, there are $|P : P'| - |P : P_1P'| \geq p - 1$ ways to

choose λ . Hence, by reciprocity, $T_1^0(G, p') \geq p - 1$, and the lemma is proved. ■

LEMMA 6. *If $P < H < G$ then*

$$T_1^0(H, p') \leq T_1^0(G, p').$$

Proof. Take $\lambda \in \text{Irr}_1^0(H, p')$. As $p \nmid \lambda^G(1)$ there exists $\chi \in \text{Irr}(\lambda^G)$ with $p \nmid \chi(1)$. Since

$$H \cap \ker \chi = \ker \chi_H \leq \ker \lambda,$$

by reciprocity the group

$$H/\ker \chi_H = H/H \cap \ker \chi \cong H \ker \chi / \ker \chi$$

is not p -nilpotent because some of its epimorphic image is the non- p -nilpotent group $H/\ker \lambda$. Then $G/\ker \chi$ is not p -nilpotent: it contains the non- p -nilpotent subgroup $H \ker \chi / \ker \chi$. Thus χ is contained in $\text{Irr}_1^0(G, p')$ and the result follows by reciprocity. ■

LEMMA 7. *If G is not p -nilpotent then $T_1(G, P, p') \geq |P : P'| - p^\alpha$, where p^α is the order of a Sylow p -subgroup of G/G' .*

Proof. Let H/G' be a normal p -complement of G/G' . Then $|G : H| = p^\alpha$ and H is not p -nilpotent. Thus [Is, Theorem 6.31] $P_1 = H \cap P$ is not contained in P' , and there is $\lambda \in \text{Lin}(P)$ such that P_1 is not contained in $\ker \lambda$. The number of such λ is equal to

$$|P : P'| - |P : P_1 P'| = |P : P'| - p^\alpha.$$

As above, there exists $\chi \in \text{Irr}(\lambda^G) \cap \text{Irr}_1(G, P, p')$, and the result follows by reciprocity. ■

Remark. Lemma 7 generalizes Lemma 3(a).

LEMMA 8. *If P is cyclic and G is not p -nilpotent then $P \leq G'$.*

Proof. This is a corollary of Tate's theorem [Is, Theorem 6.31]. However, we deduce the lemma from more elementary results. Suppose that P is not contained in G' . We may assume that $P > P \cap G' > 1$. Because $P \cap G' \cap Z(G) = 1$ we have $P \cap Z(G) = 1$. Take in G/G' a normal subgroup H/P' of index p . Set $F = N_G(P)$ and take T , a p' -Hall subgroup of F (Schur–Zassenhaus). Now $F \cap H$ is a normal subgroup of index p in F and $T \leq F \cap H$. By the Frattini Lemma and Schur–

Zassenhaus one obtains

$$F = (F \cap H)N_F(T) = (P \cap H)TN_F((T) = (P \cap H)N_F(T).$$

Since $N_F(T) = T \times P_1$, where $1 < P_1 \leq P$, $C_F(P_1) \geq \langle P, T \rangle = F$. By the Burnside normal p -complement theorem F is not p -nilpotent. So $P_1 = 1$ since by the above $p \nmid |Z(F)|$, a contradiction. ■

A group G with $T_1(G, p') = 0$ deserves special investigation. Put $\text{Irr}(G, p') = \text{Irr}_1(G, p') \cup \text{Lin}(G)$.

PROPOSITION 9. *Suppose that a group G satisfies $T_1(G, p') = 0$. If $P \in \text{Syl}_p(G)$, $N = N_G(P)$ then the following assertions hold:*

- (a) $N \cap G' = P'$.
- (b) $N = P \times A$, where A is abelian.
- (c) $G = AL$, a semi-direct product; here $L = P^G$ is the normal closure of P in G ,
- (d) $N_L(P) = P$.

Proof. Take $\lambda \in \text{Irr}(N, p')$. Because $p \nmid \lambda^G(1)$ there exists λ^0 in $\text{Irr}(\lambda^G) \cap \text{Irr}(G, p')$. By the condition $\lambda^0(1) = 1$. In particular $T_1(N, p') = 0$. Now $D = N \cap (\cap \ker \lambda^0) = \cap \ker \lambda = N'$ (here λ runs over the set $\text{Lin}(N)$). By the above $N' \geq N \cap G' \geq N'$ so $N' = N \cap G'$.

Using this reasoning we may take instead of N any subgroup H of G such that $P \leq H$. So $P' = P \cap G'$. Because $P' \leq P \cap N' \leq P \cap G' = P'$ one has $P' = P \cap N'$. By what has been proved N/P is abelian. Hence $N' \leq P$ and $P' = P \cap N' = N'$. Thus $P' = N' = N \cap G'$ and (a) is proved.

Now $P' \leq \Phi(P) \leq \Phi(N)$. Since $N/P' = N/N'$ is abelian, N is nilpotent $[G]$, $N = P \times A$. Now A is abelian, and (b) is proved.

Let $L = P^G$, the normal closure of P in G . Then $G = LN$ (Frattini). Set $M = N \cap L = N_L(P)$ and assume that $P < M$. Take $\lambda \in \text{Irr}^{\#}(M/P) = \text{Lin}^{\#}(M/P)$. Since $p \nmid \lambda^G(1)$ there exists $\lambda^0 \in \text{Irr}(\lambda^G) \cap \text{Irr}(G, p')$ and $\lambda^0(1) = 1$ by the condition. Thus $\lambda_M^0 = \lambda$, $M \cap \ker \lambda^0 = \ker \lambda \geq P$; hence $L \leq \ker \lambda^0$ for any choice of λ . Since M is not contained in $\ker \lambda$, M is not contained in $\ker \lambda^0$. Moreover L is not contained in $\ker \lambda^0$, a contradiction. Thus $M = N \cap L = P$ and (d) is proved. In particular $A \cap L = 1$. Now $G = LN = LPA = LA$ implies $G = AL = AP^G$, a semi-direct product. The proposition is proved. ■

Remark 1. If a group $G = PH$ is a semi-direct product with $P \in \text{Syl}_p(G)$ is such that $N_G(P) = P$ then as known G is solvable (Mann and Shalev [MaS] also reported to me the proof of this result). This proof uses the classification of finite simple groups. Hence Proposition 9 shows that G with $T_1(G, p') = 0$ is solvable. In fact, by Proposition 9, P coincides

with its normalizer in P^G . Since P^G is p -nilpotent (Thompson) it is solvable. Now G is solvable since G/P^G is abelian (Proposition 9).

Remark 2. Let $G(p') = \cap \ker \chi$, where χ runs over the set $\text{Irr}_1(G, p')$. It is well known [B1] that $G(p')$ is p -nilpotent. Now we prove that $\text{Irr}_1(PG(p'), p')$ is empty for $P \in \text{Syl}_p(G)$, i.e., $G(p')$ is solvable (Remark 1). Take $\lambda \in \text{Irr}_1(PG(p'), p')$, $\chi \in \text{Irr}(\lambda^G)$. Because $G(p')$ is not contained in $\ker \chi$ and χ is not linear, $p \mid \chi(1)$. Hence p divides the degrees of all irreducible constituents of λ^G . Therefore

$$|G : PG(p')| \lambda(1) = \lambda^G(1) \equiv 0 \pmod{p}.$$

Since $p \nmid |G : PG(p')|$, $p \nmid \lambda(1)$. Thus $\text{Irr}_1(PG(p'), p')$ is empty and $PG(p')$ is solvable by the previous remark. In particular $G(p')$ is solvable.

Remark 3. Let $G(p) = \cap \ker \chi$, where χ runs over the set $\text{Irr}_1(G(p)) = \{\tau \in \text{Irr}(G) \mid p \mid \tau(1)\}$. If $\lambda \in \text{Irr}_1(G(p))$, $\chi \in \text{Irr}(\lambda^G)$ then $p \nmid \lambda(1)$ since $G(p)$ is not contained in $\ker \chi$. Therefore $p \nmid \lambda(1)$. Thus one has

$$\text{Irr}_1(G(p)) = \text{Irr}_1(G(p), p').$$

As Michler [Mic] showed a Sylow p -subgroup of $G(p)$ is abelian and normal (but it is possible for $G(p)$ to be non-solvable).

In the sequel $\varphi(*)$ denotes the Euler number theoretic function.

COROLLARY. *If G is not p -nilpotent then*

$$T_1(G, P, p') \geq \varphi(|P : P'|).$$

Proof. This follows from Lemma 3(a). ■

LEMMA 10. *Suppose that G is not p -nilpotent and that P is abelian. If $T_1(G, P, p') = \varphi(|P : P'|)$ then the following assertions hold:*

- (a) $N_{G^*P}(P) = P$.
- (b) $N_G(P) = (C(s), C(p)) \times P_2$, where $|P : P_2| = p$.
- (c) $G/G' \cong C(s) \times P_2$.

Proof. Set $N = N_G(P)$, $N_1 = N_{G^*P}(P) = N \cap G^*P$. Since $G = (G^*P)N = G^*N$, $G/G' \cong N/N \cap G'$, and (c) follows from (a) and (b) in view of the transfer theorem. It remains to prove (a) and (b).

By the Burnside normal p -complement theorem N is not p -nilpotent; in particular $P < N$. Hence by Lemmas 3 and 4, and the corollary of Lemma 3, one obtains

$$\varphi(|P|) \leq T_1(N, P, p') \leq T_1(G, P, p') = \varphi(|P|),$$

and $T_1(N, P, p') = T_1(G, P, p') = \varphi(|P|)$. Thus by Lemma 3(b) one has

$$N = (C(s), C(p)) \times P_2, \quad \text{where } s|(p-1), |P:P_2| = p.$$

Hence (b) is true.

Let

$$N = LP \ (L \cap P = 1), \ N_1 = L_1P, \ L_1 \leq L, \ |L| = s, \ |L_1| = s_1.$$

Then $s_1|s$. Suppose that $s_1 > 1$, i.e., $P < N_1$.

Let μ be the sum of all linear constituents of $(1_p)^G$. Obviously $\text{Irr}(\mu) = \text{Lin}(G/G'P)$. Since $G/G'P \cong N/N_1$ one has $\mu(1) = |N:N_1| = s/s_1$. Then we have (Sylow)

$$((1_p)^G - \mu)(1) = |G:P| - s/s_1 \equiv s - s/s_1 \not\equiv 0 \pmod{p}.$$

Hence there exists $\chi \in \text{Irr}((1_p)^G) \cap \text{Irr}_1(G, p')$.

Suppose that $P \leq \ker \chi$. Let $D = P^G$ be the normal closure of P in G . By the Frattini lemma $DN = G$, so that $G/D \cong N/N \cap D$. Now $N/N \cap D$ is abelian since $N' \leq P \leq N \cap D$ by (b). Hence $G' \leq D$ and $G' \leq \ker \chi$, a contradiction since χ is not linear. Thus $\chi \in \text{Irr}_1(G, P, p')$. Since $1_p \notin \tau_1(N, P, p')_p$ (this follows from (b)), by reciprocity

$$1 + \varphi(|P|) \leq T_1(G, P, p'),$$

a contradiction. Hence $s_1 = 1$, i.e., $N_1 = P$. Since $N > P$, $s > 1$ and the lemma is proved. ■

Note that PG' from Lemma 10 is p -nilpotent (the Burnside normal p -complement theorem) and solvable (Remark 1 after Proposition 9). In particular G is a solvable group of p -length 1.

2

In this section we prove the first main theorem of this paper.

THEOREM A. *Suppose that G is not p -nilpotent and*

$$T_1(G, P, p') = \varphi(|P:P'|).$$

Then the following assertions hold:

(a) G contains a normal subgroup K such that $K \cap P = P'$, $K \leq G'$. In particular [Is, Theorem 6.31] the subgroup K is p -nilpotent.

(b) $\bar{G} = G/K$ is not p -nilpotent.

(c) $G/G'P \cong C(s)$, $s > 1$, $s | (p-1)$.

(d) $\bar{N} = N_G(P)K/K = C(s)\bar{P} = (C(s), C(p)) \times \bar{P}_2$, $|\bar{P} : \bar{P}_2| = p$.

(e) $G/G' \cong C(s) \times P_2$, where P_2 is a p -group.

(f) \bar{P} coincides with its normalizer in $\bar{P}\bar{G}'$.

Proof. Let D be the intersection of kernels of those characters $\lambda \in \text{Lin}(P)$ for which $\text{Irr}(\lambda^G) \cap \text{Lin}(G)$ is non-empty. Take for each such λ a character $\lambda^0 \in \text{Irr}(\lambda^G) \cap \text{Lin}(G)$. Then $\lambda_p^0 = \lambda$.

Take $\mu \in \text{Lin}(P) - \text{Lin}(P/D)$. Then $\text{Irr}(\mu^G) \cap \text{Lin}(G)$ is empty. Now since $p \nmid \mu^G(1)$ there exists $\chi \in \text{Irr}(\mu^G) \cap \text{Irr}_1(G, p')$. By reciprocity $\chi \in \text{Irr}_1(G, P, p')$. Set

$$\tau^0 = \sum \lambda^0, \quad \text{where } \lambda \text{ runs over the set } \text{Lin}(P/D).$$

Then, by reciprocity,

$$\text{Lin}(P) \subseteq \text{Irr}((\tau^0 + \tau_1(G, P, p'))_P).$$

By the above $T_1(G, P, p') = \varphi(|P : P'|)$ implies that $|P : D| = p$, $\langle \chi_P, \psi_P \rangle = 0$, for distinct $\chi, \psi \in \text{Irr}_1(G, P, p')$, and χ_P is multiplicity free. Set $\sigma = \tau^0 + \tau_1(G, P, p')$. Since $\text{Irr}(\sigma_P) = \text{Lin}(P)$, $K = \ker \sigma$ satisfies $K \cap P = P'$, so K is p -nilpotent [Is, Theorem 6.31]. Take $\nu \in \text{Lin}(G)$. If $\chi \in \text{Irr}_1(G, P, p')$ then $\nu\chi \in \text{Irr}_1(G, p')$. Let $\chi_P = \lambda_1 + \dots + \lambda_d$, where $\text{Irr}(\chi_P) = \{\lambda_1, \dots, \lambda_d\} \subseteq \text{Lin}(P)$. Suppose that $P \leq \ker \nu\chi$. Then $(\nu\chi)_P = d \cdot 1_P$. Hence $\nu_P \lambda_i = 1_P$ for all i , and $\lambda_1 = \dots = \lambda_d = \bar{\nu}_P$, a contradiction since $d = \chi(1) > 1$. Thus $\nu\chi \in \text{Irr}_1(G, P, p')$ so that $\nu\tau_1(G, P, p') = \tau_1(G, P, p')$. Take $x \in G - G'$ and suppose that $\nu \in \text{Lin}(G)$ is such that $\nu(x) \neq 1$. Then $\tau_1(G, P, p')(x) = 0$ and $K \leq \ker(\tau_1(G, P, p')) \leq G'$.

Let us show that G/K is not p -nilpotent. One has $K = P'K_0$, where K_0 is a normal p -complement of K . Obviously G/K_0 is not p -nilpotent, so we may assume $K_0 = 1$. Then $K = P' \leq \Phi(G)$ and G/K is not p -nilpotent by the well-known property of the Frattini subgroup $[G]$.

Put $\bar{G} = G/K$. Then $\varphi(|\bar{P}|) = \varphi(|P : P'|)$ and

$$\varphi(|\bar{P}|) \leq T_1(\bar{G}, \bar{P}, p') \leq T_1(G, P, p') = \varphi(|P : P'|) = \varphi(|\bar{P}|);$$

so $T_1(\bar{G}, \bar{P}, p') = \varphi(|\bar{P}|)$ and $\bar{P} = PK/K \cong P/P'$ is abelian. Hence the remaining assertions of the theorem follow from Lemma 10. ■

Remark. It is easy to show that the subgroup PG' from Theorem A is p -nilpotent. Hence PG' is solvable by Theorem A(f) and Remark 1 following Proposition 9. In particular, G is solvable and its p -length is equal to 1.

3

We now prove the second main theorem of this paper.

THEOREM B. $T_1^0(G, p') = p - 1$ if and only if

$$G = (C(s), C(p)), \quad \text{or} \\ G = (C(s), C(p))M = C(s)(C(p), M),$$

semi-direct products with the kernels $M, (C(p), M)$, respectively.

Proof. Let M be a maximal normal subgroup of G such that G/M is not p -nilpotent. Lemma 5 implies

$$p - 1 = T_1^0(G, p') \geq T_1^0(G/M, p') \geq p - 1$$

so $T_1^0(G, p') = p - 1 = T_1(G/M, p')$. Therefore M is defined uniquely.

(i) $M = 1$.

Then any epimorphic image of G is p -nilpotent, so G contains only one minimal normal subgroup R . Because G/R is p -nilpotent, $p \mid |R|$.

(1i) $p \nmid |G : R|$.

Then $P \leq R$ and $T_1^0(G, p') = T_1(G, P, p')$. Hence, by Lemma 3(a), one has

$$p - 1 \geq T_1(G, P, p') \geq \varphi(|P : P'|), \quad |P : P'| = p, \\ |P| = p, G = (C(s), C(p))$$

(the last equality follows from Theorem A). In the sequel let $p \mid |G : R|$.

(2i) R is a p -subgroup.

Then $G = HR$, where H is maximal in G (since R is not contained in $\Phi(G)[G]$). If $P_1 \in \text{Syl}_p(H)$ is normal in H , then $N_G(P_1) \geq \langle H, N_R(P_1) \rangle > H$, and $P_1 > 1$ is normal in G , a contradiction. Obviously $C_G(R) = R$.

Let L/R be a normal p -complement of G/R . Then L is not p -nilpotent. Take $\lambda \in \text{Lin}(P) - \text{Lin}(P/R)$ (R is not contained in P' by [G]), $\chi \in \text{Irr}(\lambda^G)$. As above, $R \leq G'$. Then G' is not contained in $\ker \chi$ (since R is not contained in $\ker \chi$), so χ is not linear, and, obviously, $G/\ker \chi = G$

is not p -nilpotent. Since $p \nmid \lambda^G(1)$, we may take χ so that $p \nmid \chi(1)$. Hence $\chi \in \text{Irr}_1^0(G, p')$. By reciprocity

$$|\text{Lin}(P) - \text{Lin}(P/R)| = |P : P'| - |P : P'R| \leq T_1^0(G, p').$$

Since P is not cyclic, $|P| \geq p^2$. Now $RP' < P$ and $|P : P'| - |P : P'R| \geq \varphi(|P : P'|) > p - 1$, a contradiction.

(3i) R is not p -solvable.

Then R is not p -nilpotent so $P_1 = P \cap R$ is not contained in P' [Is, Theorem 6.31]. Take

$$\lambda \in \text{Lin}(P) - \text{Lin}(P/P_1), \quad \chi \in \text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p').$$

Then $R < G'$, $\chi \in \text{Irr}_1^0(G, p')$, and by reciprocity

$$T_1^0(G, p') \geq |\text{Lin}(P) - \text{Lin}(P/P_1)| = |P : P'| - |P : P'P_1| > p - 1,$$

a contradiction.

(ii) $M > 1$.

Then $G/M \cong (C(s), C(p))$ by (i). As in the proof of Theorem A, we may show that $M = \ker(\tau_1^0(G, p')) < G'$; the equality $M = \ker \tau_1^0(G, p')$ follows from

$$T_1^0(G/M, p') = p - 1 = T_1^0(G, p').$$

Thus $|G' : M| = p$, $G/G' \cong C(s)$. It remains to prove that $G' = (C(p), M)$.

Take $\chi \in \text{Irr}(G, P, p')$. Since $p \nmid |G/\ker \chi|$ and $p \nmid |G : G'|$, $G/\ker \chi$ is not p -nilpotent, i.e., $\chi \in \text{Irr}_1^0(G, p')$. Therefore

$$\text{Irr}_1(G, P, p') \subseteq \text{Irr}_1^0(G, p').$$

Since the opposite inclusion is evident, one has

$$T_1(G, P, p') = T_1^0(G, p') = p - 1.$$

Then Theorem A implies $|P| = p$ and $N_{G'}(P) = P$ so $G' = (C(p), M)$. Since a group G with such a structure satisfies the condition of the theorem, the proof is finished. ■

In the same way, using Lemma 8, we may prove the following:

PROPOSITION 11. *Suppose that G is not p -nilpotent and $P \in \text{Syl}_p(G)$ is cyclic. Then the following conditions are equivalent:*

- (a) $T_1^0(G, p') = |P| - 1$.
- (b) $G = (C(s), P)M$, a semi-direct product with normal subgroup M , $N_{PM}(P) = P$.

Obviously G from Proposition 11 is solvable.

4

Let

$$\begin{aligned} \text{Irr}_1^1(G, p') &= \{\chi \in \text{Irr}_1(G, p') \mid G/\ker \chi \text{ is not } p\text{-solvable}\}, \\ \tau_1^1(G, p') &= \sum \chi, \quad \text{where } \chi \text{ runs over the set } \text{Irr}_1^1(G, p'); \\ T_1^1(G, p') &= \tau_1^1(G, p')(1). \end{aligned}$$

In this section we give a lower estimate of $T_1^1(G, p')$ for a non- p -solvable G with a cyclic Sylow p -subgroup P .

LEMMA 12. *Let a non- p -nilpotent group $G = AP$, where $P \cong C(p^m)$, $P \in \text{Syl}_p(G)$ is normal in G , $A \cap P = 1$. If $T_1(G, P, p') \leq |P| + 2$ then one of the following assertions holds:*

- (a) $p = 3, G \cong D(12)$, a dihedral group of order 12.
- (b) $p = 3, G = C(4)C(3)$.
- (c) $G = (C(s), P), s \mid (p - 1)$.

Proof. Set $L = C_A(P)$. Then $G/L \cong (C(s), P), s \mid (p - 1)$. If $L = 1$ one obtains a group (c). Assume that $L > 1$. Obviously $p \neq 2$. We have $T_1(G/L, PL/L, p') = |P| - 1$ (see [Is, Theorem 6.34]). Let $\mu \in \text{Irr}^*(L), \lambda \in \text{Lin}(P)$ with $\ker \lambda = 1$, and $\chi \in \text{Irr}((\mu \times \lambda)^G)$. Since $P \leq G'$ (Lemma 8) and $\ker \chi \cap P = 1, \chi(1) > 1$. Since $\text{Irr}_1(G) = \text{Irr}_1(G, p'), \chi \in \text{Irr}_1(G, P, p')$. Hence

$$T_1(G, P, p') \geq T_1(G/L, PL/L, p') + \chi(1) = |P| - 1 + \chi(1).$$

By condition $\chi(1) \leq 3$. If there is a character $\tau \in \text{Irr}_1(G, P, p') - \text{Irr}(G/L, PL/L, p') - \{\chi\}$ we obtain

$$T_1(G, P, p') \geq |P| - 1 + \chi(1) + \tau(1) > |P| + 2,$$

a contradiction. Thus

$$\text{Irr}_1(G, P, p') = \text{Irr}_1(G/L, PL/L, p') \cup \{\chi\}, \chi(1) \leq 3.$$

Then all characters from $\text{Irr}^*(L)$ are G -conjugate. This implies that L is an elementary abelian q -group, a prime $q \neq p$. Moreover L is a minimal normal subgroup of G . Suppose that $|P| > p, P_1$ is a subgroup of order p in P . Take in $\text{Lin}(P/P_1)$ a faithful character λ_1 , and take $\mu \in \text{Irr}^*(L)$. If $\tau \in \text{Irr}((\mu \times \lambda_1)^G)$ then $\tau \neq \chi$ and $\tau \in \text{Irr}_1(G, P, p')$, a contradiction. Thus $|P| = p$. Suppose that L is not contained in G' ; then $L = Z(G)$ and

$|L| = 2$ (since all characters from $\text{Lin}^*(L)$ are G -conjugate). Hence $G = D(4p)$, a dihedral group of order $4p$, or $G = C(4)P$. Then $T_1(G, P, p') = 2(p-1) \leq p+2$, so $p = 3$ and $G \cong D(12)$ or $G = C(4)C(3)$.

Suppose that $L \leq G'$. Then $G' = P \times L$ and

$$|G| - |G|/|P| - |G|/|L| + |G|/|P||L| = \chi(1)^2 \in \{4, 9\}.$$

Hence $s|L|p - sp - s|L| + s = t \in \{4, 9\}$.

(i) Suppose that $t = 4$. Then $s|4$. If $s = 4$ then $|L|p = |L| + p$ and $p = 2$, a contradiction. Let $s = 2$. Then $|L|p = |L| + p + 1$, a contradiction.

(ii) Now let $t = 9$. Then $s|9$. If $t = 9$ then $|L|p = p + |L|$ and $p = 2$, a contradiction. If $s = 3$ then $|L|p = p + |L| + 2$, which is impossible. The lemma is proved. ■

THEOREM 13. *Let G be a non- p -solvable group with cyclic Sylow p -subgroup P . Then*

- (a) $T_1^1(G, p') \geq |P| + 1$.
- (b) If $p > 3$ then $T_1^1(G, p') \geq |P| + 2$.

Proof. Let G be a counterexample of minimal order. Then

- (i) $O_p(G) = 1$.

Let $R > 1$ be a normal subgroup of G such that P is not contained in R . Then $R \cap P \leq \Phi(P)$, and hence R has a normal p -complement $O_p(R)$ ($\leq O_p(G) = 1$) ([Is, Theorem 6.31]). Thus $R \leq P$. By the transfer theorem $P \cap C_G(R)' \cap Z(C_G(R)) = 1$ or $R \cap C_G(R)' = 1$. Therefore $C_G(R)' \leq O_p(C_G(R))$. Since $C_G(R)$ is normal in G ,

$$O_p(C_G(R)) \leq O_p(G) = 1 \quad \text{and} \quad C_G(R) = P.$$

Since $G/C_G(R)$ is isomorphic to a subgroup of $C(p-1)$, G is solvable, a contradiction. Thus

(ii) Any non-trivial normal subgroup of G contains P and, in particular, it is not p -solvable. Moreover G contains only one minimal normal subgroup R . Since $P < R$, G/R is a p' -group. Now $R \leq G'$ by Lemma 8.

(i)–(ii) imply

$$(iii) \text{ Irr}_1(G, P, p') = \text{Irr}_1^1(G, p'), T_1(G, P, p') = T_1^1(G, p').$$

Set $N = N_G(P)$, $N_1 = N \cap R (= N_R(P))$.

By Burnside's normal p -complement theorem one has

- (iv) N and N_1 are not p -nilpotent.

$$(v) T_1(G, P, p') \geq T_1(N, P, p').$$

For the proof see Lemma 4(a).

Now suppose that $R < G$.

By the assumption on G and (v) it follows that $T_1(N, P, p') \leq |P| + 2$. By virtue of Lemma 12 we know the structure of N . By the Frattini lemma one has $G = RN$ so that $R = G'$. Now Lemma 12 and (v) imply

(vi) If N is not a Frobenius group then $N \cong D(12)$, $N_1 \cong D(6)$.

Suppose that $N \cong D(12)$, $N_1 \cong D(6)$. Then $|P| = 3$ and $C_R(P) = P$. So by the Feit–Thompson theorem one has $R \in \{A_5, PSL(2, 7)\}$ [FT]. Hence, by (i), one has $G \in \{S_5, PGL(2, 7)\}$. Since

$$\begin{aligned} T_1(S_5, P, 3') &= 18 > 5 = |P| + 2, \\ T_1(PGL(2, 7), P, 3') &= 30 > 5 = |P| + 2, \end{aligned}$$

we obtain a contradiction. Thus

(vii) If $R < G$ then $N = (C(s), P)$, $N_1 = (C(s_1), P)$, $s_1 | s$, $s_1 < s$.

We continue to consider the case $R < G$.

Take $\lambda \in \text{Lin}(N) - \text{Lin}(N/N_1)$. Since $\lambda^G(1) = |G : N| \equiv 1 \pmod{p}$ (Sylow) and $G' = R$ is not contained in $\ker \lambda$ there exists (by reciprocity) a $\chi \in \text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')$. Now

$$|\text{Lin}(N)| - |\text{Lin}(N/N_1)| = s - s/s_1.$$

Thus

$$T_1(G, P, p') \geq T_1(N, P, p') + s - s/s_1 = |P| - 1 + s - s/s_1.$$

In particular $T_1(G, P, p') \geq |P| + 1$, and (a) is proved in the case $R < G$. Next we assume that $p > 3$, $s - s/s_1 = 2$. Then

$$s = 4, s_1 = 2, N = (C(4), P), 4 | (p - 1).$$

Let $\varphi \in \text{Irr}_1(N)$, $\chi \in \text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')$. Then $\chi \in \text{Irr}_1(G, P, p')$. If for some choice of φ one has $|\text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')| > 1$ then $\varphi(1) = 4$ implies (by reciprocity)

$$T_1(G, P, p') \geq T_1(N, P, p') + \varphi(1) \geq |P| - 1 + 4 = |P| + 3,$$

and the theorem is proved. Thus we may assume that for any $\varphi \in \text{Irr}_1(N)$ one has $|\text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')| = 1$. Now for χ from $\text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')$ one has $\langle \varphi^G, \chi \rangle = 1$ (since in the contrary case the contribution of φ in $T_1(G, P, p')$ is at least 8, and we obtain $T_1(G, P, p') \geq |P| - 1 + 4 = |P| + 3$). In particular for our χ one has $\chi(1) \equiv \varphi^G(1) \equiv 4 \pmod{p}$.

Take $\tau \in \text{Irr}_1(G, P, p') = \text{Irr}_1(G, p')$. Then τ is faithful. Hence τ_N has a non-linear irreducible constituent ψ . By the above, in view of $\tau \in \text{Irr}(\psi^G) \cap \text{Irr}_1(G, P, p')$, it follows that $\tau(1) \equiv 4 \pmod{p}$.

Take $\lambda \in \text{Lin}(N) - \text{Lin}(N/N_1)$. Then $\lambda^G(1) = |G : N| \equiv 1 \pmod{p}$ and $\text{Irr}(\lambda^G) \subseteq \text{Irr}_1(G)$ (since any irreducible constituent of λ^G does not contain G' in its kernel). Hence there exists $\chi \in \text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')$. If the contribution of λ in $T_1(G, p')$ is at least 2 then $T_1(G, p') \geq |P| - 1 + s - s/s_1 + 1 \geq |P| + 2$. Hence we may assume that $\langle \lambda^G, \chi \rangle = 1$ for any $\chi \in \text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')$. By the proved $\chi(1) \equiv 4 \pmod{p}$. Since $p > 3$ there exists $\tau \in \text{Irr}((\lambda)^G - \chi) \cap \text{Irr}_1(G, p')$ (because $p \nmid (\lambda^G - \chi)(1)$) so that the contribution of λ in $T_1(G, p')$ is greater than 1, a contradiction. Thus

(viii) $G' = R = G$; i.e., $s = s_1$, $N = N_1$, G is a non-abelian simple group.

Suppose that N is not a Frobenius group. By Lemma 12 one has $N = D(12)$ or $N = C(4)C(3)$, $|P| = 3$. Then $T_1(N, 3') = 4 = |P| + 1$, and (a) is proved by virtue of (v). Therefore in the sequel we assume that $p > 3$. Then $N = (C(s), P)$ by (v) and Lemma 12.

As above it is easy to prove the following assertions:

(ix) If $\varphi \in \text{Irr}_1(N)$ then $|\text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')| = 1$. Moreover if $\chi \in \text{Irr}(\varphi^G) \cap \text{Irr}_1(G, p')$ then $\langle \varphi^G, \chi \rangle = 1$.

(x) $\chi(1) \equiv s \pmod{p}$ for all $\chi \in \text{Irr}_1(P, p') = \text{Irr}_1(G, p')$.

(xi) If $\lambda \in \text{Lin}^*(N)$ then by virtue of (x) one has $|\text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')| > 1$ since $\lambda^G(1) \equiv 1 \pmod{p}$, $\text{Irr}(\lambda^G) \subseteq \text{Irr}_1(G)$. Hence

(xii) $T_1(G, P, p') \geq T_1(N, p') + 2(s - 1) = |P| - 1 + 2(s - 1)$.

Since G is a counterexample,

(xiii) $s = 2$.

If, in (xi), we have $|\text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')| > 2$ then we obtain in (xii) strong inequality, so $T_1(G, P, p') > |P| - 1 + 2 = |P| + 1$, a contradiction (since G is a counterexample). Now $\chi(1) \equiv 2 \pmod{p}$ for all $\chi \in \text{Irr}_1(G, p')$ by (x).

Let $\lambda \in \text{Lin}^*(N)$. Because $\text{Irr}(\lambda^G) \subseteq \text{Irr}_1(G)$, $\lambda^G(1) \equiv 1 \pmod{p}$, (x) implies that $|\text{Irr}(\lambda^G) \cap \text{Irr}_1(G, p')| \geq (p + 1)/2 > 2$, a contradiction with the above. Then theorem is proved. ■

Conjectures. 1. Suppose that G and P are as in Theorem 13. Then

$$T_1^1(G, p') \geq |P| + p.$$

2. If G is a non- p -solvable group then $T_1^1(G, p') \geq 2p$.

QUESTION. Suppose that $P \in \text{Syl}_p(G)$ and $P < H < G$. Classify all pairs $H < G$ such that $T_1(H, P, p') = T_1(G, P, p')$.

5

In this section we consider groups G with $|\text{Irr}_1(G, p')| = 1$.

PROPOSITION 14. *Suppose that p is a prime divisor of $|G|$, $|\text{Irr}_1(G, p')| = 1$, $\text{Irr}_1(G, p') = \{\chi\}$, $\ker \chi = 1$. Then the following assertions hold:*

- (a) G' is the only minimal normal subgroup of G .
- (b) $p \nmid |G : G'|$.
- (c) If G is p -solvable then $G = (C(p^n - 1), E(p^n))$.
- (d) If $p = 2$ then $G' < G$.

(e) If G is not p -solvable, then G' is simple and all characters from $\text{Irr}_1(G', p')$ are conjugate under G .

Proof. Suppose that $p \mid |G : G'|$. Then G contains a normal subgroup H of index p . By Clifford's theory $\chi_H = \eta \in \text{Irr}(H)$, and

$$\eta^G = \chi_1 + \cdots + \chi_p,$$

where χ_1, \dots, χ_p are distinct irreducible characters of G of p' -degree $\chi(1)$, a contradiction. Hence (b) is proved.

Let N be a minimal normal subgroup of G . By the condition the degrees of all non-linear irreducible characters of G/N are divisible by p , so G/N has a normal p -complement H/N (Thompson) and (b) implies $H = G$ whence G/N is a p' -number. Since $N > \ker \chi = 1$ all irreducible characters of G/N are linear by the Ito theorem [Is], Theorem 6.15]. Hence G/N is abelian, $N = G'$, and (a) is proved.

By (b) one has $P \leq G'$ (here $P \in \text{Syl}_p(G)$). Suppose G is p -solvable. Then $P = G'$ is a minimal normal subgroup of G by (a). Hence $\text{Irr}_1(G) = \text{Irr}_1(G, p')$ ([Is, Theorem 6.15]) and $|\text{Irr}_1(G)| = 1$. Therefore $G = (C(p^m - 1), E(p^m))$ by [S] and (c) is proved.

Suppose that $p = 2$ and $G' = G$. Since all characters from $\text{Irr}(G) - \{1_G, \chi\}$ have even degrees, $|G| \equiv 1 + \chi(1)^2 \equiv 2 \pmod{4}$. Therefore G contains a normal 2-complement, contradicting (b).

Suppose that G is not p -solvable. Let $\chi_{G'} = e(\lambda_1 + \cdots + \lambda_t)$ be the Clifford decomposition. Since G' is a direct product of isomorphic non-abelian simple groups, $\lambda_1, \dots, \lambda_t$ are non-linear. Suppose that $\lambda \in \text{Irr}_1(G', p') - \{\lambda_1, \dots, \lambda_t\}$. Then by (b) and the Clifford theory one has $\text{Irr}(\lambda^G) \subseteq \text{Irr}_1(G, p')$. Since $\chi \notin \text{Irr}(\lambda^G)$ one obtains a contradiction. Thus we have $\text{Irr}_1(G', p') = \{\lambda_1, \dots, \lambda_t\}$, and $\lambda_1, \dots, \lambda_t$ are conjugate under G . In particular $\lambda_1(1) = \dots = \lambda_t(1)$. Hence G' is not a direct product of two or more non-abelian simple groups. This proves (e). The proposition is proved. ■

Remark 1. If $\text{Irr}_1(G, p') = \{\chi\}$ then $\ker \chi$ is p -nilpotent and solvable (see Remark 1 after Proposition 9).

Remark 2. Put $c(G, p') = |\text{Irr}_1(G, p') \cup \text{Lin}(G)|$. Alperin and McKay conjectured that $c(G, p') = c(N_G(P), p')$ (see [Mic], p. 227). If a non- p -solvable group G satisfies $|\text{Irr}_1(G, p')| = 1$ and $G/\ker \chi$ satisfies the Alperin–McKay conjecture (we do not assume here that this conjecture holds in general) then it is easy to prove the following assertions:

- (a) $p < 5$.
- (b) $N_G(P)/P' = (C(p^d - 1), E(p^d))$, where $|P : P'| = p^d$. Moreover $N_G(P) = P, P' = \Phi(P)$.
- (c) $G/G' \cong C(p^d)$.

Conjecture. If $|\text{Irr}_1(G, p')| = 1$ then G is p -solvable.

6

In this section we prove the following

PROPOSITION 15. *Let $P \in \text{Syl}_p(G)$. Then the following assertions are equivalent:*

- (a) $T_1(G, p') = 0$.
- (b) $T_1(G, P, p') = 0$.

Proof. Obviously $T_1(G, p') = 0$ implies $T_1(G, P, p') = 0$.

Assume that (b) is true but (a) is not true. Let $P^G = L$ be the normal closure of P in G and H the normal p -complement of G (note that G is p -nilpotent by Lemma 7). Then $N = N_G(P) = P \times A$. Suppose that A is not abelian. Take $\mu \in \text{Irr}_1(A)$ and $\lambda \in \text{Lin}^\#(P)$. Since $p \nmid (\lambda \times \mu)^G(1)$ there exists $\chi \in \text{Irr}((\lambda \times \mu)^G)$ such that $p \nmid \chi(1)$. Now $\chi \in \text{Irr}_1(G, P, p')$ by reciprocity, a contradiction. Thus A is abelian. By the Frattini lemma one has $G = LN = LA$. Suppose that $\tau \in \text{Irr}_1(G, p')$. By the condition one has $P \leq \ker \tau$. Hence $P^G = L \leq \ker \tau$, a contradiction, since G/L is abelian and τ is non-linear. Thus (a) is true and the proposition is proved. ■

Let $G = P \times A$, where $P \in \text{Syl}_p(G)$ and A is not abelian. Then $T_1(G, p') > T_1(G, P, p')$.

PROPOSITION 16. *Let $P \in \text{Syl}_p(G), L = P^G$. Then the following assertions are equivalent:*

- (a) $T_1(G, p') = 0$.
- (b) $T_1(L, p') = 0$ and $G' = L'$.

Proof. Suppose that (a) is true. Then $T_1(L, p') = 0$ by Lemma 4(b). To prove $G' = L'$ we may assume that $L' = 1$. Then G is abelian by Proposition 9 and $G' = 1$. Thus (b) is proved. Suppose that (b) is true but (a) is not true. Take $\chi \in \text{Irr}_1(G, p')$. By (b) and the Clifford theorem one has $\text{Irr}(\chi_L) \subseteq \text{Lin}(L)$. Hence by (b) we have $G' = L' \leq \ker \chi$ and χ is linear, a contradiction. ■

ACKNOWLEDGMENTS

I am indebted to Avinoam Mann, Aner Shalev, and Pavel Shumjatsky for useful discussions.

REFERENCES

- [B1] YA. G. BERKOVICH, Degrees of irreducible characters and normal p -complements, *Proc. Amer. Math. Soc.* **106** (1989), 33–35.
- [BM] YA. BERKOVICH AND A. MANN, On sums of degrees of irreducible characters, in “Structure Theory of Set Addition,” pp. 197–219, CIRM, Marseille, 1993.
- [Br] A. BRANDIS, Über die Grade irreduziblen Charaktere endlichen Gruppen, *J. Algebra* **97** (1985), 23–29.
- [FT] W. FEIT AND J. G. THOMPSON, Finite groups which contain a self-centralizing subgroups of order 3, *Nagoya Math. J.* **21** (1962), 185–197.
- [G] W. GASCHÜTZ, Über die Φ -Untergruppen endlicher Gruppen, *Math. Z.* **58** (1953), 160–170.
- [Is] I. M. ISAACS, “Character Theory of Finite Groups,” Academic Press, New York, 1976.
- [K] G. KARPILOVSKY, “Group Representations,” Vol. 1, part. B, North-Holland, Amsterdam, 1992.
- [MaS] A. MANN AND A. SHALEV, private communication.
- [Mic] G. O. MICHLER, Modular representation theory and the classification of finite simple groups, *Proc. Symp. Pure Math.* **47** (1987), 223–232.
- [S] G. SEITZ, Finite groups having only one irreducible representation of degree greater than one, *Proc. Amer. Math. Soc.* **19** (1968), 459–461.